

ON THE MAXIMAL RANK IN A SUBSPACE OF MATRICES

by ROY MESHULAM

[Received 4th July 1984]

On the maximal rank in a subspace of matrices

LET $M_n(F)$ be the space of $n \times n$ matrices over a field F , and let W be a linear subspace of $M_n(F)$.

Flanders [2] proved that if $\dim W > rn$ and $|F| \geq r+1$, then W contains a matrix of rank $> r$. He also characterized the subspaces W such that $\dim W = rn$ and W contains no matrix of rank $> r$.

In this note we prove a lower bound on the maximal rank attained in a subspace of matrices (Theorem 1). We then use this bound to derive Flanders' results (Theorems 2 and 3) without restrictions on F .

Let $[n]$ denote $\{1, \dots, n\}$, and let $<$ be the lexicographic order on $[n] \times [n]$. ($(i, j) < (i_1, j_1)$ iff $i < i_1$ or $i = i_1$ and $j < j_1$.)

For $A \in M_n(F)$ denote by $p(A) \in [n] \times [n]$, the location of A 's lexicographically first non-zero entry:

$$p(A) = \min \{(i, j) : A(i, j) \neq 0\}$$

For a collection $\mathcal{A} = \{A_1, \dots, A_m\}$ of $n \times n$ matrices, construct an $n \times n$ matrix B as follows: $B(k, l) = 1$ if $(k, l) = p(A_i)$ for some $1 \leq i \leq m$, and $B(k, l) = 0$ otherwise.

Denote by $\rho(\mathcal{A})$ the minimal number of lines in B (a line is either a row or a column) which cover all 1's in B .

THEOREM 1. *Let $\mathcal{A} = \{A_1, \dots, A_m\} \subset M_n(F)$. Then span \mathcal{A} contains a matrix of rank $\geq \rho(\mathcal{A})$.*

Proof. Call a set of entries in a matrix independent, if it contains no two entries on the same line. By König's Theorem ([4], Theorem 5.1.4 in [3]), the maximal size of an independent set of 1's in a 0-1 matrix, is equal to the minimal number of lines, which cover all 1's in that matrix. Hence if $\rho(\mathcal{A}) = r$, then there exist $1 \leq i_1, \dots, i_r \leq m$ such that $\{p(A_{i_j}) : 1 \leq j \leq r\}$ is independent.

Let $p(A_{i_j}) = (k_j, l_j)$ for $1 \leq j \leq r$, then $S = \{k_1, \dots, k_r\}$ and $T = \{l_1, \dots, l_r\}$ are both of cardinality r . For $1 \leq j \leq r$ define $B_j = A_{i_j}[S | T] \in M_r(F)$ (the minor determined by restricting the entries to $S \times T$).

We shall prove the theorem by showing that span $\{B_1, \dots, B_r\}$ contains a non-singular matrix.

We may assume that $k_1 < \dots < k_r$. Let h be the permutation on $[r]$ for which: $l_{h(1)} < \dots < l_{h(r)}$. Denote the j th row of B_j by b_j .

Clearly B_j 's first $(j-1)$ rows are zero, $b_j(k)=0$ for $1 \leq k < h^{-1}(j)$, and $b_j(h^{-1}(j)) \neq 0$. Let C be the $r \times r$ matrix, whose rows are b_1, \dots, b_r . C is non-singular, because by the preceding remarks, permuting C 's rows according to h , we obtain an upper triangular matrix, with non-zeros on the diagonal. Let $D_j = B_j C^{-1}$ for $1 \leq j \leq r$. It is easy to check that the following holds:

$$\begin{aligned} \text{For all } 1 \leq j \leq r: D_j \text{'s first } j-1 \text{ rows are zero} \\ \text{and } D_j \text{'s } j\text{th row is the } j\text{th unit vector.} \end{aligned} \tag{1}$$

Claim 1. If D_1, \dots, D_r satisfy (1), then there exists a 0-1 combination of D_1, \dots, D_r which is non-singular.

Proof. We use induction on r . The case $r=1$ is trivial. Assume $r > 1$. For $1 \leq j \leq r-1$ let $D'_j = D_j([r-1] | [r-1]) \in M_{r-1}(F)$. D'_1, \dots, D'_{r-1} satisfy (1) for $r-1$, and so, by induction there exist $x_1, \dots, x_{r-1} \in \{0, 1\}$ such that $\sum_{j=1}^{r-1} x_j D'_j$ is non-singular.

Now, since $D_r(i, j) = 0$ for all $(i, j) \neq (r, r)$, and $D_r(r, r) = 1$, we obtain by expanding the bottom row:

$$\det \left(\sum_{j=1}^{r-1} x_j D_j + D_r \right) = \det \left(\sum_{j=1}^{r-1} x_j D'_j \right) + \det \left(\sum_{j=1}^{r-1} x_j D'_j \right) \tag{2}$$

But $\det(\sum_{j=1}^{r-1} x_j D'_j) \neq 0$, so (2) implies that $\sum_{j=1}^{r-1} x_j D_j$ and $\sum_{j=1}^{r-1} x_j D_j + D_r$ cannot both be singular. ■

We return to the proof of the theorem. By the claim $\sum_{j=1}^r x_j D_j$ is non-singular for some x_j 's, and therefore $\sum_{j=1}^r x_j B_j = (\sum_{j=1}^r x_j D_j)C$ is also non-singular. This implies that $\text{rank}(\sum_{j=1}^r x_j A_{ij}) \geq r$. ■

The next result had been proved by Flanders [2], for $|F| \geq r+1$:

THEOREM 2. If W is a subspace of $M_n(F)$, and $\dim W > m$, then W contains a matrix of rank $> r$. ■

Proof. Choose a basis $\mathcal{A} = \{A_1, \dots, A_t\}$ of W . By performing a gaussian elimination on $\{A_1, \dots, A_t\}$ (regarding them as n^2 dimensional vectors), we may assume that $p(A_1), \dots, p(A_t)$ are all distinct. Since a line in a matrix covers n entries, we cannot cover $p(A_1), \dots, p(A_t)$ by less than t/n lines. Therefore $\rho(\mathcal{A}) \geq t/n > r$, which by Theorem 1 implies that $W = \text{span } \mathcal{A}$ contains a matrix of rank $> r$. ■

Next we discuss a certain extremal case of Theorem 2.

Let F^n be the space of n -tuples over F . Denote by $x \otimes y \in M_n(F)$, the Kronecker product of $x, y \in F^n$. For $A, B \subset F^n$, let $A \otimes B = \text{span } \{x \otimes y : x \in A, y \in B\}$.

The following result had been proved by Flanders [2], under the assumptions $|F| \geq r+1$ and $\text{char}(F) \neq 2$. Atkinson and Lloyd [1] had obtained it assuming only $|F| \geq r+1$.

THEOREM 3. *Suppose $W \subset M_n(F)$ is a subspace of dimension m , such that for all $A \in W$, $\text{rank}(A) \leq r$. Then either $W = E \otimes F^n$ or $W = F^n \otimes E$, for some r dimensional subspace $E \subset F^n$.*

Proof. Let $\mathcal{A} = \{A_1, \dots, A_m\}$ be a basis of W . As in Theorem 2, we may assume that $p(A_1), \dots, p(A_m)$ are all distinct. W does not contain a matrix of rank $> r$, therefore by Theorem 1, $\rho(\mathcal{A}) \leq r$. Choose r lines which cover $p(A_1), \dots, p(A_m)$. Since each line covers at most n of the $p(A_i)$'s, it follows that the lines are pairwise disjoint, and that each of them consists entirely of $p(A_i)$'s.

Hence, either all r lines are rows, or all r lines are columns.

We shall assume the first case—that is: $p(A_1), \dots, p(A_m)$ form r rows. (The case of columns is treated similarly).

Next we note that if $Q_1, Q_2 \in M_n(F)$ are non-singular, then the maximal rank in $Q_1 W Q_2$ is equal to the maximal rank in W , and $W = E_1 \otimes E_2$ for some $E_1, E_2 \subset F^n$ iff $Q_1 W Q_2 = (Q_1 E_1) \otimes (E_2 Q_2)$.

In particular, by performing the same row permutation on all matrices in W , we may assume that $p(A_1), \dots, p(A_m)$ consist of the first r rows.

Clearly, by gaussian elimination on A_1, \dots, A_m (regarded as vectors in F^n), we may obtain a new basis $\{B_{ij} : 1 \leq i \leq r, 1 \leq j \leq n\}$ of W , such that $B_{ij}(i, j) = 1$ and $B_{ij}(k, l) = 0$ for all $1 \leq k \leq r, 1 \leq l \leq n$ such that $(k, l) \neq (i, j)$.

Claim 2. B_{ij} is zero, except for the j th column.

Proof. We have to show that $B_{ij}(k, l) = 0$ for $l \neq j$ and $r + 1 \leq k \leq n$ (for $1 \leq k \leq r$ this is known). Since our claim is invariant under row and column permutations, it suffices to prove it for specific i, j, k, l (which satisfy $l \neq j$ and $r + 1 \leq k \leq n$), say $i = j = r, k = l = r + 1$. That is, we show that $B_r(r + 1, r + 1) = 0$. let $C_{ij} = B_{ij}([r + 1] | [r + 1]) \in M_{r+1}(F)$, and define $E_{ij} \in M_r(F)$ for $1 \leq i, j \leq r$ by $E_{ij}(k, l) = \delta_{ik} \delta_{jl}$.

Let $P \subset [r] \times [r]$. As $C_p(i, r + 1) = 0$ for all $p \in P, 1 \leq i \leq r$, we have:

$$\det \left(\sum_{p \in P} C_p \right) = \det \left(\sum_{p \in P} E_p \right) \left(\sum_{p \in P} C_p(r + 1, r + 1) \right). \tag{3}$$

Since W does not contain a matrix of rank $> r$, it follows that $\det(\sum_{p \in P} C_p) = 0$, and so if $P \subset [r] \times [r]$ satisfies:

$$\det \left(\sum_{p \in P} E_p \right) \neq 0 \tag{4}$$

Then $\sum_{p \in P} C_p(r + 1, r + 1) = 0$.

It is clear that the sets $P = \{(1, 1), (2, 2), \dots, (r - 2, r - 2), (r - 1, r), (r, r - 1)\}$ ($P = \{(1, 1)\}$ for $r = 1$), and $P_1 = P \cup \{(r, r)\}$, both satisfy (4),

and so:

$$\sum_{p \in P} C_p(r+1, r+1) = \sum_{p \in P_1} C_p(r+1, r+1) = 0.$$

This implies $B_r(r+1, r+1) = C_r(r+1, r+1) = 0$. ■

We complete the proof of Theorem 3, by showing that for every $1 \leq i \leq r$ there exists $x_i \in F^n$, such that for every $1 \leq j \leq n$ $B_{ij} = x_i \otimes e_j$ (e_j is the j th unit vector in F^n).

In view of Claim 2, we only have to show that for $1 \leq j_1, j_2 \leq n$, the j_1 th column of B_{i_1} is equal to the j_2 th column of B_{i_2} . Again by permuting rows and columns it suffices to prove (for example) that $B_{11}(r+1, 1) = B_{12}(r+1, 2)$. Using the notations of Claim 2, let

$$C = C_{11} + C_{12} + (C_{23} + C_{34} + \dots + C_{r+1})$$

By Claim 2: $C(r+1, 1) = B_{11}(r+1, 1)$, $C(r+1, 2) = B_{12}(r+1, 2)$. C , being an $(r+1) \times (r+1)$ minor of a matrix in W is singular, because W has no member of rank $> r$. On the other hand it is clear that:

$$\det(C) = (-1)^r (C(r+1, 1) - C(r+1, 2))$$

Therefore $C(r+1, 1) = C(r+1, 2)$ and so: $B_{11}(r+1, 1) = B_{12}(r+1, 2)$. ■

Remark. Atkinson and Lloyd [1] have extended Flanders' classification, by proving that if $W \subset M_n(F)$ does not contain a matrix of rank $> r$, $\dim W \geq rn - r + 1$ and $|F| \geq r + 1$, then W is r -decomposable (that is: $W \subset E_1 \otimes F^n + F^n \otimes E_2$ for some subspaces $E_1, E_2 \subset F^n$ such that $\dim E_1 + \dim E_2 = r$).

Contrary to Theorems 2 and 3, this result does depend on the field, as the following example, which has been suggested by the referee, indicates: Let W be the 5-dimensional space of all

$$\begin{pmatrix} a & 0 & 0 \\ c & b & 0 \\ d & e & a+b \end{pmatrix} \tag{5}$$

over $GF(2)$. Clearly W does not contain a non-singular matrix, yet W is not 2-decomposable. For otherwise W' —the space of all matrices of the form (5) over say, $GF(4)$ —would also be 2-decomposable, which is impossible since W' contains non-singular matrices.

Acknowledgment

I would like to thank Nathan Linial and Michael Rabin for their help.

REFERENCES

1. M. D. Atkinson and S. Lloyd, 'Large subspaces of matrices of bounded rank', *Quarterly J. Math.* 31 (1980). 253-262.

2. H. Flanders, 'On spaces of linear transformations with bounded rank', *J. London Math. Soc.* 37 (1962), 10–16.
3. M. Hall, Jr., *Combinatorial Theory*, Blaisdell, Waltham, Mass. 1967.
4. D. König, 'Graphok és matrixok', *Mat. Fiz. Lapok* 38 (1931), 115–119.

*Institute of Mathematics,
Hebrew University
Jerusalem 91904
Israel*