

# COLLAPSING WALLS THEOREM

IGOR PAK\* AND ROM PINCHASI†

ABSTRACT. Let  $P \subset \mathbb{R}^3$  be a pyramid with the base a convex polygon  $Q$ . We show that when other faces are *collapsed* (rotated around the edges onto the plane spanned by  $Q$ ), they cover the whole  $Q$ .

## 1. INTRODUCTION

Let  $P$  be a convex pyramid in  $\mathbb{R}^3$  over the base  $Q$ , which is a convex polygon in a horizontal plane. Think of the other faces  $F$  of  $P$  as the “walls” of a wooden box, and that each wall  $F$  is hinged to the base  $Q$  along the edge. Suppose now that the walls are “collapsed”, i.e. rotated around the edges towards the base onto the horizontal plane. The question is: *do they cover the whole base  $Q$ ?*



FIGURE 1. An impossible configuration of four collapsing walls of a pyramid leaving a hole in the base.

At first, this may seem obvious, but in fact the problem is already non-trivial even in the case of four-sided pyramids, which can possibly have some obtuse dihedral angles (see Figure 1). Formally, we have the following result:

**Collapsing Walls Theorem.** *Let  $P \subset \mathbb{R}^3$  be a pyramid over a convex polygon  $Q$ . For a face  $F$  of  $P$ , denote by  $e_F$  the edge between  $F$  and the base:  $e_F = F \cap Q$ , and let  $A_F$  denotes the result of rotation of  $F$  around  $e_F$  in the direction of  $P$ , onto the plane which contains  $Q$ . Then*

$$Q \subseteq \cup_F A_F,$$

where the union is over all faces  $F$  of  $P$ , different from  $Q$ .

For example, suppose pyramid  $P$  in the theorem has a very large height, so that all walls are nearly vertical. The theorem then implies that every point  $z \in Q$  has an orthogonal projection into the interior of some edge  $e$  of  $Q$ . This is a classical result

---

*Date:* 28 March 2009.

\*School of Mathematics, University of Minnesota, Minneapolis, MN; Email: pak@umn.edu.

†Department of Mathematics, Technion, Haifa, Israel; Email: room@math.technion.ac.il.

with a number of far-reaching generalizations (see [Pak, §9]). Thus, the collapsing walls theorem can be viewed as yet another generalization of this result (cf. Section 3).

## 2. PROOF OF THE THEOREM

Consider  $\mathbb{R}^3$  endowed with the standard Cartesian coordinates  $(x_1, x_2, x_3)$ . Without loss of generality assume that the plane  $H$  spanned by  $Q$  is horizontal, i.e. given by  $x_3 = 0$ , and that  $P$  is contained in the half-space  $x_3 \geq 0$ . Denote by  $F_1, \dots, F_m$  the faces of  $P$  different from  $Q$ , by  $H_i$  the planes spanned by  $F_i$ , and by  $e_i = F_i \cap Q$  the edges of  $Q$ , for all  $1 \leq i \leq m$ .

Denote by  $\Phi_i$  the rotation about  $e_i$  of  $H_i$  onto  $H$  (the rotation is performed in the direction dictated by  $P$ , so that throughout the rotation  $H_i$  intersects the interior of  $P$ ). Similarly, let  $A_i = \Phi_i(F_i)$  is the rotation of the face  $F$  of  $P$  onto  $Q$ ,  $1 \leq i \leq m$ . We need to show that every point in  $Q$  lies in  $\cup_{i=1}^m A_i$ . Without loss of generality we can take this point to be the origin  $O$ .

Further, denote by  $L_i = H_i \cap H$  the line through  $e_i$ . Let  $r_i$  be the distance from the origin to  $L_i$ , and let  $\alpha_i$  be the dihedral angle of  $P$  at  $e_i$ , i.e the angle between  $H$  and  $H_i$  which contains  $P$ .

Suppose now  $F_1$  is a face such that

$$\tau_i = r_i \cdot \tan \frac{\alpha_i}{2} \quad \text{is minimized at } \tau_1.$$

We will show that the origin  $O$  is contained in  $A_1$ . In other words, we prove that if  $O \notin A_1$ , then  $\tau_i < \tau_1$  for some  $i > 1$ .

Let  $z \in H_1$  such that the rotation of  $z$  onto  $Q$  is the origin:  $\Phi_1(z) = O$ . It suffices to show that  $z \in F_1$ . Let  $\mathbf{v} = (v_1, v_2, 0)$  be the unit vector that is a normal to  $L_1$  in the horizontal plane. It is easy to see that

$$\overrightarrow{Oz} = (r_1(1 - \cos \alpha_1)v_1, r_1(1 - \cos \alpha_1)v_2, r_1 \sin \alpha_1).$$

To prove the theorem, assume to the contrary that  $z \notin F_1$ . Then there exists a face of  $P$ , say  $F_2$ , such that  $H_2$  separates  $z$  from the origin. Denote by  $y$  the closest point to  $z$  on  $L_2$ , and by  $\alpha'$  the angle between the line  $(zy)$  and the horizontal plane  $H$ , where the angle is taken with the half-plane of  $H$  which contains  $Q$  (and thus the origin). In this notation, the above condition implies that  $\alpha' > \alpha_2$ .

Without loss of generality we may assume that line  $L_2$  is given by equations  $x_2 = r_2$  and  $x_3 = 0$ . Then

$$y = (r_1(1 - \cos \alpha_1)v_1, r_2, 0),$$

and

$$\cos \alpha' = \cos \widehat{Oyz} = \frac{r_2 - r_1(1 - \cos \alpha_1)v_2}{\sqrt{r_1^2 \sin^2 \alpha_1 + (r_2 - r_1(1 - \cos \alpha_1)v_2)^2}}.$$

Note that the function  $x/\sqrt{a^2 + x^2}$  is monotone increasing as a function of  $x$ , and that  $v_2 \leq 1$ . We get

$$\cos \alpha' \geq \frac{r_2 - r_1(1 - \cos \alpha_1)}{\sqrt{r_1^2 \sin^2 \alpha_1 + (r_2 - r_1(1 - \cos \alpha_1))^2}}.$$

Applying  $\cos \alpha' < \cos \alpha_2$ , we conclude:

$$(1) \quad \frac{r_2 - r_1(1 - \cos \alpha_1)}{\sqrt{r_1^2 \sin^2 \alpha_1 + (r_2 - r_1(1 - \cos \alpha_1))^2}} < \cos \alpha_2.$$

Recall the assumption that  $\tau_1 \leq \tau_2$ . This gives  $r_1 \tan \frac{\alpha_1}{2} \leq r_2 \tan \frac{\alpha_2}{2}$ , or

$$(2) \quad \frac{r_2}{r_1} \geq \frac{\tan \frac{\alpha_1}{2}}{\tan \frac{\alpha_2}{2}}.$$

The rest of this section is dedicated to showing that both (1) and (2) are impossible. This gives a contradiction with our assumptions and proves the claim. We split the proof into two cases depending on whether the dihedral angle  $\alpha_2$  is acute or obtuse. In each case we repeatedly rewrite (1) and (2), eventually leading to a contradiction.

**Case 1** (obtuse angles). Suppose  $\frac{\pi}{2} < \alpha_2 < \pi$ . In this case  $\cos \alpha_2 < 0$ , and (1) is equivalent to

$$(3) \quad 1 + \frac{r_1^2 \sin^2 \alpha_1}{(r_2 - r_1(1 - \cos \alpha_1))^2} < \frac{1}{\cos^2 \alpha_2},$$

and

$$(4) \quad \frac{r_1 \sin \alpha_1}{r_2 - r_1(1 - \cos \alpha_1)} > \tan \alpha_2.$$

This can be further rewritten as:

$$(5) \quad \frac{r_2}{r_1} < 1 - \cos \alpha_1 + \frac{\sin \alpha_1}{\tan \alpha_2}.$$

Now (5) and (2) together imply

$$\frac{\tan \frac{\alpha_1}{2}}{\tan \frac{\alpha_2}{2}} < 1 - \cos \alpha_1 + \frac{\sin \alpha_1}{\tan \alpha_2},$$

which is impossible. Indeed, suppose for some  $0 < a, b < \pi$ , we have

$$(6) \quad \frac{\tan \frac{a}{2}}{\tan \frac{b}{2}} < 1 - \cos a + \frac{\sin a}{\tan b}.$$

Dividing both sides by  $(\tan \frac{a}{2})$ , after some easy manipulations, we conclude that (6) is equivalent to

$$(7) \quad \frac{1}{\tan \frac{b}{2}} < \sin a + \frac{1 + \cos a}{\tan b},$$

which in turn is equivalent to

$$(8) \quad \left( \frac{1}{\tan \frac{b}{2}} - \frac{1}{\tan b} \right) \sin b < \cos(a - b).$$

Since the left hand side of (8) is equal to 1, we get a contradiction and complete the proof in Case 1.

**Case 2** (right and acute angles). Suppose now that  $0 < \alpha_2 \leq \frac{\pi}{2}$ . Then  $\cos \alpha_2 \geq 0$ , and  $0 < \tan \frac{\alpha_2}{2} \leq 1$ . Let us first show that the numerator of (1) is nonnegative, i.e. that  $r_2 \geq r_1(1 - \cos \alpha_1)$ . From the contrary assumption we have  $r_2/r_1 < (1 - \cos \alpha_1)$ . Together with (2), this implies:

$$1 - \cos \alpha_1 > \frac{r_2}{r_1} \geq \frac{\tan \frac{\alpha_1}{2}}{\tan \frac{\alpha_2}{2}} \geq \tan \frac{\alpha_1}{2},$$

which is impossible for all  $0 < \alpha_1 < \pi$ .

From above, we can exclude the right angle case  $\alpha_2 = \frac{\pi}{2}$ , for else the l.h.s. of (1) is nonnegative, while r.h.s. is equal to zero. Thus,  $\cos \alpha_2 > 0$ . Therefore, the inequality (1) in this case can be rewritten as

$$(9) \quad 1 + \frac{r_1^2 \sin^2 \alpha_1}{(r_2 - r_1(1 - \cos \alpha_1))^2} > \frac{1}{\cos^2 \alpha_2},$$

and

$$(10) \quad \frac{r_1 \sin \alpha_1}{r_2 - r_1(1 - \cos \alpha_1)} > \tan \alpha_2.$$

Note now that (10) coincides with (4). Since (6) holds for all  $0 < a, b < \pi$ , we obtain the contradiction verbatim the proof in Case 1. This completes the analysis of Case 2 and finishes the proof of the theorem.  $\square$

### 3. FINAL REMARKS

**3.1.** The collapsing walls theorem extends verbatim to higher dimensions. Moreover, it also extends to every polytope  $P \subset \mathbb{R}^d$ , as follows. Fix one facet  $Q$  of  $P$  and assume all other facets  $F$  of  $P$  are rotated around the affine subspace  $H_F \cap H$  onto the hyperplane  $H$  containing  $Q$ , then they cover the whole facet  $Q$ . Here  $H_F$  denotes the hyperplane that contains the facet  $F$ . We refer to [PP], where this result is proved in full generality, and is used to show that a smaller polyhedron can always be sequentially cut out of a bigger polyhedron, in any dimension.

**3.2.** Let us note that when the walls of a pyramid are collapsed *outside*, rather than *onto* the base, they are pairwise non-intersecting (see Figure 2). We leave this easy exercise to the reader.

**3.3.** Continuing with the example of “vertical walls” as given in the introduction right after the theorem, recall that for the center of mass  $z = \text{cm}(Q)$ , there are at least two such edges onto which orthogonal projection of  $z$  lies in the interior (see e.g. [Pak, §9]).<sup>1</sup> It would be interesting to see if this result extends to the setting of the theorem (of course, the notion of the center of mass would have to be modified appropriately). Let us note here that the center of mass result is closely related to the four vertex theorem [Tab], and fails in higher dimension [CGG].

---

<sup>1</sup>One can give a construction with there is only one such edge, if the center of mass is replaced by a general point in  $Q$  (see [CGG] and [Pak, §9]).




FIGURE 2. Walls of a pyramid collapsing outside the base do not intersect.

**3.4.** The proof of the theorem is based on an implicit subdivision of  $Q$  given by the smallest of the linear functions  $\tau_i$  at every point  $z \in Q$ . Recall that  $\tau_i$  is a weighted distance to the edge  $e_i$ . Thus this subdivision is in fact a weighted analogue of the dual Voronoi subdivision in the plane (see [Aur, For]). As a consequence, computing this subdivision can be done efficiently, both theoretically and practically.

**Acknowledgments.** The authors are thankful to Yuri Rabinovich for the interest in the problem. The first author was partially supported by the National Security Agency and the National Science Foundation. The second author was supported by the Israeli Science Foundation (grant No. 938/06).

#### REFERENCES

- [Aur] F. Aurenhammer, Voronoi diagrams—a survey of a fundamental geometric data structure, *ACM Comput. Surv.* **23** (1991), 345–405.
- [CGG] J. H. Conway, M. Goldberg and R. K. Guy, Problem 66-12, *SIAM Review* **11** (1969), 78–82.
- [For] S. Fortune, Voronoi diagrams and Delaunay triangulations, in *Computing in Euclidean geometry* (F. Hwang and D. Z. Du, eds.), World Scientific, Singapore, 1995, 225–265.
- [Pak] I. Pak, *Lectures on Discrete and Polyhedral Geometry*, monograph to appear, 2009; available at <http://www.math.umn.edu/~pak>
- [PP] I. Pak and R. Pinchasi, How to cut out a convex polyhedron, preprint (2009).
- [Tab] S. Tabachnikov, Around four vertices, *Russian Math. Surveys* **45** (1990), 229–230.