

On the number of distinct directions of planes determined by n points in \mathbb{R}^3

Rom Pinchasi*

August 27, 2007

Abstract

We show that any set of n points in \mathbb{R}^3 , that is not contained in a plane, determines at least $\lfloor n/2 \rfloor + 1$ two-dimensional affine hyper-planes, no two of which are parallel.

1 Introduction

In 1970, Scott [S70] raised the following questions: What is the minimum number of different directions assumed by the connecting lines of (1) n points in the plane, not all on a line, (2) n points in 3-space, not all on a plane?

In 1982, Ungar [U82] solved the first problem, by verifying Scott's conjecture that in the plane the above minimum is equal to $2\lfloor n/2 \rfloor$, for any $n > 3$. Ungar's proof is a real gem, a brilliant application of the method of *allowable sequences* invented by Goodman and Pollack [GP81], [GP93].

Scott's conjecture regarding a set of n points in \mathbb{R}^3 was recently settled in [PPS07] as follows:

Theorem 1 ([PPS07]). *Every set of $n \geq 6$ points in \mathbb{R}^3 , not all of which are on a plane, determines at least $2n - 5$ different directions if n is odd, and at least $2n - 7$ different directions if n is even. This bound is sharp for every odd $n \geq 7$.*

In this paper we consider another natural analogue of Scott's conjecture in \mathbb{R}^2 to three dimensions. Namely, we consider distinct directions of two-dimensional planes determined by n points in \mathbb{R}^3 .

Let P be a finite set of points in \mathbb{R}^3 . A (affine) plane H is said to be *determined* by P if it passes through (at least) three non-collinear points in P . Similarly, we say that a line ℓ is

*Mathematics Dept., Technion—Israel Institute of Technology, Haifa 32000, Israel.
room@math.technion.ac.il.

determined by P if ℓ passes through (at least) two points of P . A straight line segment e is said to be determined by P , if e is delimited by two points in P .

We consider all the planes determined by a set P of n points in \mathbb{R}^3 and we wish to bound from below the number of distinct directions of these planes in terms of n . Here, two planes are of the same direction if they are parallel in the affine space \mathbb{R}^3 , or in other words if the normal vectors to these planes are the same.

Clearly, if the set P is coplanar, then it determines at most one plane and hence we will assume that the set P is not contained in a plane. By considering small examples it is natural to conjecture the following:

Conjecture 1. *Let P be a set of n points in \mathbb{R}^3 that is not contained in a plane. Then P determines at least n planes with pairwise distinct directions.*

If one cares only about distinct planes determined by P , regardless of their affine position, then it was shown by Hanani ([H54]) that a non-coplanar set of n points in \mathbb{R}^3 determines at least n distinct planes. For every $n \geq 4$ it is not hard to come up with a construction of n points that determine precisely n planes with pairwise distinct directions. One can take for example $n - 2$ points on one line ℓ and two points on another line that is not coplanar with ℓ . Then the resulting set of n points determines only n planes, with pairwise distinct direction.

In Section 3, we make a first step towards giving an affirmative answer to Conjecture 1. We prove the following partial result:

Theorem 2. *Let P be a set of n points in \mathbb{R}^3 that is not contained in a plane. Then P determines at least $\lfloor n/2 \rfloor + 1$ planes, no two of which are parallel.*

2 Some preliminary notions and results

A generalization and strengthening of the result in [U82] about the minimum number of distinct directions of lines determined by n points in the plane, is given in [PPS03]. In order to introduce this result we need the following definition that will be important also throughout the rest of the paper.

Definition 1. Two segments in \mathbb{R}^2 are called *convergent* if they are opposite edges in a convex quadrilateral, or if they are collinear. See Figure 1

In other words, two non-collinear segments are not convergent if and only if a line through one of the segments meets the other segment. The notion of convergent segments in the plane was first introduced by Kupitz (see [K84]), who conjectured that a geometric graph on n vertices with no pair of convergent edges has at most $2n - 2$ edges. Kupitz conjecture was proved by Katchalski, Last, and Valtr in [KL98] and [V98]. This result is in fact tight for $n \geq 4$, by a construction of Kupitz ([K84]).




Figure 1: Two convergent segment

In [PPS03] it is shown that any set P of n non-collinear points in the plane determines at least $2\lfloor n/2 \rfloor$ segments, no two of which are convergent. This result clearly implies the result in [U82], as two parallel segments are always convergent.

The result in [PPS03] follows very easily if P is a set of points no three of which are collinear. In fact, under this condition P determines n segments no two of which are convergent. The interesting and more difficult cases are when more than two points in P may be collinear.

In this respect the result in [KL98] and [V98] is formulated for sets of points that are in general position, i.e., contain no three points on a line. However, it is an easy consequence that the same result is true for any set of points. As we will need this more general version, we give a short argument showing how it can be deduced from the result in [KL98] and [V98]:

Theorem 3 (Simple corollary of [KL98] and [V98]). *Let P be a set of n points in the plane and let E be a collection of segments determined by P . If no two segments in E are convergent, then E consists of at most $2n - 2$ segments.*

Proof. Recall that two segments are not convergent exactly when a line through one segment meets the other segment. Hence, if two segments e and f are not convergent, and e' and f' are two segments containing e and f , respectively, then e' and f' are not convergent. Therefore, we can assume that all the segments in E are maximal segments determined by P . That is, each segment e in E is delimited by the two extreme points of P on the line containing e .

Consider now a small perturbation of a point $x \in P$. By a small perturbation we refer to a continuous perturbation that does not cause new collinearities among the points of P . This perturbation effects those segments delimited by x . We claim that a small enough perturbation cannot cause two segments in E to become convergent. Indeed, let e and f be two segments in E . The assertion is clear if e and f share a common endpoint. Therefore, assume that they do not. Since e and f are not convergent, assume without loss of generality that the line ℓ through e meets the segment f . If ℓ meets the relative interior of f , then no small perturbation of a point $x \in P$ can change this. It is left to consider the case where ℓ meets a point $y \in P$ delimiting the segment f . Because e is maximal, and e and f do not share a common endpoint, it follows that y is contained in the relative interior of e . But then the line through f meets that relative interior of e and this cannot be changed by a small perturbation of a point in P .

To complete the proof consider small perturbations of the points in P such that the resulting set of points is in general position. Then apply the theorem of Katchalski, Last, and Valtr to conclude that E consists of at most $2n - 2$ edges. ■

3 Proof of the main theorem

Lemma 1. *Let $P = \{p_1, \dots, p_n\}$ be a non-coplanar set of n points in \mathbb{R}^3 , contained in the open half-space $\{z > 0\}$. We assume also that the points of P are in general position with respect to the origin in the sense that there is no plane determined by P that passes through O . Let e_1, \dots, e_m be a set of m straight line segments determined by P . Assume that for every $1 \leq i < j \leq m$ either the plane through O and e_i meets e_j , or the plane through O and e_j meets e_i , or both. Then P determines at least $\frac{1}{2}m + 1$ planes with pairwise distinct directions.*

Remark: Note that because $P \subset \{z < 0\}$, the condition on e_1, \dots, e_m is equivalent to that the central projection through O of the segments e_1, \dots, e_m , on a plane parallel to $\{z = 0\}$, results in a collection of segments no two of which are convergent.

Proof. It will be convenient for us to assume that the set P lies not only in the half-space $\{z > 0\}$ but in fact in the open quadrant $\{y > 0, z > 0\}$. To see that there is no loss of generality in this assumption, we show that one can find a linear transformation which takes P to be in the quadrant $\{y > 0, z > 0\}$. Consider the linear transformation which takes a point $(x, y, z) \in \mathbb{R}^3$ to the point $(\epsilon x, \epsilon y, z)$, for a very small, but positive, ϵ . This mapping takes P to a set of points P' contained in a narrow cone around the positive part of the z -axis. A rotation which takes the positive part of the z -axis to the ray $\{x = y = z > 0\}$ will take P' to a set of points P'' that lies in the open quadrant $\{y > 0, z > 0\}$, as desired. Observe that applying a linear transformation to P does not effect the other conditions in Lemma 1, nor the property of two planes determined by P to have distinct directions.

We will therefore assume that $P \subset \{y > 0, z > 0\}$. Let K_1 be the plane $\{z = 1\}$. Let T denote the central projection of \mathbb{R}^3 through O on the plane K_1 . That is, T takes a point $p \in \mathbb{R}^3 \setminus \{z = 0\}$ to the intersection point of the line through O and p with the plane K_1 . The image of a point $p = (a, b, c) \in \mathbb{R}^3 \setminus \{z = 0\}$ is the point $T(p) = (\frac{a}{c}, \frac{b}{c}, 1)$.

For every $i = 1, \dots, n$, let (a_i, b_i, c_i) denote the Cartesian coordinates of the point $p_i \in P$. T takes each point $p_i \in P$ to the point $q_i = (\frac{a_i}{c_i}, \frac{b_i}{c_i}, 1)$ on K_1 . By the conditions of the lemma, the segments e_1, \dots, e_m , determined by P , are projected by T to m segments on K_1 , no two of which are convergent. We conclude that q_1, \dots, q_n determine m segments on K_1 , no two of which are convergent.

Let \mathcal{D}' be the duality on the plane K_1 which takes a point $(a, b, 1)$ to the line $\{ax + by = 1\}$ on K_1 , and which takes a line $\{ax + by = 1\}$ on K_1 to the point $(a, b, 1) \in K_1$. For every $1 \leq i \leq n$, \mathcal{D}' takes the point q_i to the line $\ell_i = \{a_i x + b_i y = c_i\}$ on K_1 . Let \mathcal{L} denote the collection of these n lines on K_1 .

Observe that the y -coordinate of each point q_i is positive. This is by the definition of T

and because of the assumption that $P \subset \{y > 0, z > 0\}$. Let ℓ_i and ℓ_j be two lines in \mathcal{L} . Because the y -coordinates of both q_i and q_j are positive, it follows that \mathcal{D}' takes the segment delimited by q_i and q_j to the set of lines passing through the intersection point of ℓ_i and ℓ_j with a slope that lies in the range between the slope of ℓ_i and the slope of ℓ_j . We call the union of this set of lines the *double-wedge* determined by ℓ_i and ℓ_j . The intersection point of ℓ_i and ℓ_j is called the *apex* of the double-wedge (see Figure 2). Because the points q_1, \dots, q_n determine m segments no two of which are convergent, then the line arrangement \mathcal{L} contains m double-wedges such that among each two, there is one double wedge that contains the apex of the other double-wedge. We say that two such double-wedges are *not convergent*. See Figure 2.




Figure 2: On the left: a double wedge. On the right: two double wedges that are not convergent.

Consider now again the original set P in \mathbb{R}^3 . Let \mathcal{D} be the duality which takes a point $(a, b, c) \in \mathbb{R}^3 \setminus \{O\}$ to the plane $\{ax + by + cz = 1\}$ and which takes a plane $\{ax + by + cz = 1\}$ to the point $(a, b, c) \in \mathbb{R}^3$. This duality respects incidence relations and sends parallel planes to points collinear with O . Let H_1, \dots, H_n denote the duals of p_1, \dots, p_n , respectively, under the duality \mathcal{D} .

For every $i = 1, \dots, n$, we have $p_i = (a_i, b_i, c_i)$ and therefore, $\mathcal{D}(p_i) = H_i = \{a_i x + b_i y + c_i z = 1\}$. Let K denote the plane $\{z = z_0\}$ for large enough number z_0 , so that K lies above all the vertices in the hyper-planes arrangement \mathcal{H} consisting of H_1, \dots, H_n . The intersection of H_i with K is the line $L_i = \{a_i x + b_i y + c_i z_0 = 1\}$ on K . Let $T' : \mathbb{R}^3 \rightarrow \mathbb{R}^3$ be the linear transformation which takes a point (x, y, z) to the point $\frac{1}{z_0}(-x, -y, z)$. For each i , T' takes the line L_i to the line $\ell'_i = \{a_i x + b_i y - c_i = -\frac{1}{z_0}\}$ on the plane K_1 . Note that for z_0 large enough, this line is arbitrarily close to the line $\ell_i \in \mathcal{L}$ on the plane K_1 . Moreover, for any three indices i, j, k , the lines ℓ'_i, ℓ'_j , and ℓ'_k are concurrent whenever ℓ_i, ℓ_j , and ℓ_k are. This is easy to see because if ℓ_i, ℓ_j , and ℓ_k are concurrent, then $T(p_i), T(p_j)$, and $T(p_k)$ are collinear. But this means that p_i, p_j , and p_k are collinear, for otherwise they determine a plane passing through O , which is impossible. The collinearity of p_i, p_j , and p_k now easily implies that ℓ'_i, ℓ'_j , and ℓ'_k are concurrent. We denote by \mathcal{L}' the set of lines $\{\ell'_1, \dots, \ell'_n\}$.

We would like to show that the set P determines “many” planes with pairwise distinct directions. This is equivalent to showing that the \mathcal{D} -dual arrangement \mathcal{H} which consists of

the planes H_1, \dots, H_n determines “many” vertices, no two of which are collinear with the origin O . Because T' is a non-singular linear transformation, it is enough to show that in the arrangement $T'(\mathcal{H})$ there are “many” vertices, no two of which are collinear with the origin.

We saw already that the arrangement $T'(\mathcal{H})$ intersects K_1 in an arrangement of lines ℓ'_1, \dots, ℓ'_n , and that if z_0 is large enough, then each ℓ'_i is arbitrarily close to ℓ_i and any concurrency of lines from \mathcal{L} corresponds to concurrency of corresponding lines from \mathcal{L}' . Because \mathcal{L} determines m double-wedges no two of which are convergent, it follows that ℓ'_1, \dots, ℓ'_n as well determine m double-wedges, no two of which are convergent.

We now define a geometric graph G on the plane K_1 . The vertices of G will be vertices of the arrangement $T'(\mathcal{H})$ centrally projected on K_1 through O . Each edge of G will correspond to one of the m double-wedges, determined by ℓ'_1, \dots, ℓ'_n , no two of which are convergent. We will then show that no two edges in G are convergent. Hence, by Theorem 3, the number m of edges in G is at most $2|V(G)| - 2$, where $V(G)$ is the set of vertices of G . It will then follow that $|V(G)| \geq m/2 + 1$. This will prove the lemma because each vertex of G corresponds to a central projection of a vertex of $T'(\mathcal{H})$.

To construct G , suppose that ℓ'_i and ℓ'_j determines one of the above mentioned m double-wedges on K_1 . The apex of this double-wedge is in fact the intersection of $T'(H_i)$, $T'(H_j)$, and K_1 . Let A and B be the two extreme vertices of $T'(\mathcal{H})$ on the line of intersection of $T'(H_i)$ and $T'(H_j)$. We consider the line segment $[A, B]$, delimited by A and by B , and centrally project it on K_1 to obtain an edge of G . Hence, A and B are projected to vertices of G .

It is left to show that no two edges of G are convergent. Let e and f be two edges of G . Assume that e corresponds to the double-wedge W_{ij} determined by ℓ'_i and ℓ'_j , and that f corresponds to the double-wedge W_{kl} determined by ℓ'_k and ℓ'_l . Without loss of generality assume that W_{ij} contains the apex of the double-wedge W_{kl} . We will show that the line through e meets the edge f on K_1 , implying that e and f are not convergent.

Recall that ℓ'_i is the line $\{a_i x + b_i y - c_i = -\frac{1}{z_0}\}$ and similarly, $\ell'_j = \{a_j x + b_j y - c_j = -\frac{1}{z_0}\}$, both on the plane K_1 .

H_i is the plane $\{a_i x + b_i y + c_i z = 1\}$. As T' is the mapping defined by $T'(x, y, z) = \frac{1}{z_0}(-x, -y, z)$, $T'(H_i)$ is the plane $\{a_i x + b_i y = c_i z + \frac{1}{z_0}\}$. Similarly, $T'(H_j) = \{a_j x + b_j y = c_j z + \frac{1}{z_0}\}$. It is now easy to verify that the line of intersection of $T'(H_i)$ and $T'(H_j)$ is projected through O to the line $t_{ij} = \{(a_i - a_j)x + (b_i - b_j)y = c_i - c_j\}$ on K_1 .

Since $-\frac{a_i - a_j}{b_i - b_j}$, the slope of t_{ij} on K_1 , is never between $-\frac{a_i}{b_i}$ and $-\frac{a_j}{b_j}$, the slopes of ℓ'_i , and ℓ'_j on K_1 , respectively, it follows that t_{ij} is not contained in W_{ij} (see Figure 3).

Let R denote the plane passing through O and t_{ij} . Let R^+ denote the closed half-space bounded by R which contains the apex of W_{kl} . R^- will denote the other closed half-space bounded by R . For convenience define $T'(H_i)^+ = T'(H_i) \cap R^+$ and $T'(H_i)^- = T'(H_i) \cap R^-$. Similarly, denote $T'(H_j)^+ = T'(H_j) \cap R^+$ and $T'(H_j)^- = T'(H_j) \cap R^-$.

The apex of W_{kl} is in the convex portion of \mathbb{R}^3 bounded by $T'(H_i)^+$ and $T'(H_j)^+$.

Case 1. The apex of W_{kl} is neither on $T'(H_i)^+$ nor on $T'(H_j)^+$. Because the plane K_1




Figure 3: The projection on K_1 . A view from the origin O .

lies above all vertices of the arrangement $T'(\mathcal{H})$, then all the vertices of $T'(\mathcal{H})$ on the line of intersection of $T'(H_k)$ and $T'(H_l)$ lie on the part of this line below K_1 . Let us denote this ray by \vec{r} . Without loss of generality assume that \vec{r} crosses $T'(H_i)$ before it crosses $T'(H_j)$. It follows that \vec{r} intersects $T'(H_i)$ on $T'(H_i)^+$. The intersection point of \vec{r} and $T'(H_i)^+$ is a vertex of $T'(\mathcal{H})$ that lies in R^+ . The intersection point of \vec{r} with $T'(H_j)$ must be therefore on $T'(H_j)^-$, as \vec{r} cannot cross twice the plane $T'(H_i)$. The intersection point of \vec{r} and $T'(H_j)^-$ is a vertex of $T'(\mathcal{H})$ that lies in R^- . It follows that the edge f of G , which corresponds to the double-wedge W_{kl} must cross the line t_{ij} that contains the edge e .

Case 2. The apex of W_{kl} is either on $T'(H_i)^+$ or on $T'(H_j)^+$. Without loss of generality assume that the apex of W_{kl} is on $T'(H_i)^+$. Because K_1 lies above all vertices in the arrangement $T'(\mathcal{H})$, it follows that the line of intersection of $T'(H_k)$ and $T'(H_l)$ is contained in the plane $T'(H_i)$. Therefore, the intersection point of $T'(H_k)$, $T'(H_l)$, $T'(H_i)$, and $T'(H_j)$ is a vertex in $T'(\mathcal{H})$. The projection of this vertex through O on K_1 lies on both e and f . It follows that e and f are not convergent. ■

We are now ready to proceed to the proof of Theorem 2. As mentioned already, it is shown in [PPS03] that any non-collinear set of n points in the plane determines at least $2\lfloor n/2 \rfloor$ segments no two of which are convergent. In view of this, Theorem 2 is a corollary of Lemma 1:

Proof of Theorem 2. Translate P so that it is contained in the half-space $\{z > 0\}$ of \mathbb{R}^3 , and so that no plane determined by P is incident to the origin O . Consider now the central projection of the points of P through O on the plane $K_1 = \{z = 1\}$. The points of P are projected to n non-collinear points on K_1 . By the result in [PPS03], one can find at least $m = 2\lfloor n/2 \rfloor$ segments on K_1 determined by the images of the points of P such that no two

of these segments are convergent. The corresponding segments determined by the points of P satisfy the conditions in Lemma 1. The corollary now follows. ■

4 Suggestions and concluding remarks

Let P be a set of points in \mathbb{R}^3 , that is not contained in a plane. Call a point O *good* with respect to P if O lies outside of the convex hull of P and it is in general position with respect to P in the sense that O does not lie on any plane determined by P .

It could be tempting to conjecture that for every set P of n points in \mathbb{R}^3 , that is not contained in a plane, there is a good point O such that the central projection through O of the set P on a plane determines $2n - 2$ segments no two of which are convergent. This, together with Lemma 1, would immediately imply Conjecture 1. However, there are sets P for which such a good point O does not exist. This is strongly related to Scott's conjecture in \mathbb{R}^3 , mentioned in the introduction. Indeed, let P be a set of n points (n odd) in \mathbb{R}^3 that does not determine more than $2n - 5$ lines with pairwise distinct directions. One can take, for example, the vertices of a regular $(n - 3)$ -gon centered at the origin on the $x - y$ plane, together with the points $(0, 0, 0)$, $(0, 0, 1)$, and $(0, 0, -1)$. Any central projection of such a point set P through a good point O will determine at most $2n - 5$ segments no two of which are convergent. This is because two parallel segments will always project to two convergent segments.

It will therefore be interesting to determine the maximum number $g(n)$ such that for any set P of n non-coplanar points in \mathbb{R}^3 there exists a good point O such that the central projection of P through O on a plane determines at least $g(n)$ segments no two of which are convergent. From [PPS03] and the example above it follows that for every odd n , $n - 1 \leq g(n) \leq 2n - 5$. Similarly, one can show that for even n , $n \leq g(n) \leq 2n - 3$.

Lemma 1 implies immediately that any set P of n non-coplanar points in \mathbb{R}^3 determines at least $\lceil g(n)/2 \rceil + 1$ planes with pairwise distinct directions.

One could hope to get a good lower bound for $g(n)$ from the constructive proof of Scott's conjecture in \mathbb{R}^3 presented in [PPS07]. Yet, a close examination of the resulting set of segments with pairwise distinct directions, produced by the proof in [PPS07], does not directly imply such a lower bound.

Finally, looking into the proof of Lemma 1 one realizes that the crucial step there is proving that the edges of the geometric graph G are pairwise non-convergent. The following simple observation can be used to strengthen Lemma 1 and perhaps lead to a complete solution of Conjecture 1. The geometric graph G is constructed in such a way that the edges in G correspond to line segments determined by P , and the vertices of G correspond to planes determined by P , no two of which are parallel. In fact, if e is a segment determined by P and \tilde{e} is the corresponding edge in G , then every plane determined by P which passes through e corresponds to some point on \tilde{e} . Therefore, if e and f are two segments determined by P that happen to lie on the same plane H , then the corresponding edges \tilde{e} and \tilde{f} in G will meet at the point which corresponds to H . In particular, \tilde{e} and \tilde{f} are not convergent.

Therefore, Lemma 1 can be strengthened as follows:

Lemma 2. *Let $P = \{p_1, \dots, p_n\}$ be a non-coplanar set of n points in \mathbb{R}^3 , contained in the open half-space $\{z > 0\}$. We assume also that the points of P are in general position with respect to the origin in the sense that there is no plane determined by P that passes through O . Let e_1, \dots, e_m be a set of m straight line segments determined by P . Assume that for every $1 \leq i < j \leq m$ either the plane through O and e_i meets e_j , or the plane through O and e_j meets e_i , or e_i and e_j are coplanar. Then P determines at least $\frac{1}{2}m + 1$ planes with pairwise distinct directions.*

The following question now naturally arises: Is it true that any set of n non-coplanar points in \mathbb{R}^3 determines at least $2n - 3$ segments, every two of which are either coplanar, or their central projections through some fixed good point O are not convergent?

An affirmative answer to this question will imply that Conjecture 1 is true.

References

- [GP81] J. E. Goodman and R. Pollack, A combinatorial perspective on some problems in geometry, *Congr. Numer.* **32** (1981), 383–394.
- [GP93] J. E. Goodman and R. Pollack, Allowable sequences and order types in discrete and computational geometry, in: *New Trends in Discrete and Computational Geometry (J. Pach, ed.)*, *Algorithms Combin.* **10**, Springer, Berlin, 1993, 103–134.
- [H54] H. Hanani, On the number of lines and planes determined by d points. Technion. Israel Inst. Tech. Sci. Publ. **6** (1954/5), 58–63.
- [KL98] M. Katchalski and H. Last, On geometric graphs with no two edges in convex position, *Discrete Comput. Geom.* **19** (1998), no. 3, Special Issue, 399–404.
- [K84] Y.S. Kupitz, On pairs of disjoint segments in convex position in the plane, *Ann. Discrete Math.* **20** (1984), 203–208.
- [PPS03] J. Pach, R. Pinchasi, and M. Sharir, On the number of directions determined by a three-dimensional point set, *J. Combinat. Theory, Ser. A* **108** (2004), 1–16.
- [PPS07] J. Pach, R. Pinchasi, and M. Sharir, Solution of Scott’s Problem on the Number of Directions Determined by a Point Set in 3-Space, *Discrete Comput. Geom.* **38** (2007), 399–441.
- [S70] P. R. Scott, On the sets of directions determined by n points, *Amer. Math. Monthly* **77** (1970), 502–505.
- [U82] P. Ungar, $2N$ noncollinear points determine at least $2N$ directions, *J. Combin. Theory, Ser. A* **33** (1982), 343–347.

- [V98] P. Valtr, On geometric graphs with no k pairwise parallel edges, *Discrete Comput. Geom.* **19** (1998), no. 3, Special Issue, 461–469.